

Emulsifier for the formulation of cosmetic W/O creams and lotions

INTENDED USE

W/O emulsifier

BENEFITS AT A GLANCE

- Low usage concentration of 1.5 – 2.5%
- High compatibility with active ingredients
- Emulsifier for "light" W/O emulsions with pleasant application properties
- Emulsions with high heat and freeze stability
- Stable emulsions without co-emulsifiers and with low amounts of consistency-enhancing waxes
- Formulations with all kinds of cosmetic oils
- Liquid at room temperature

INCI (PCPC NAME)

Cetyl PEG/PPG-10/1 Dimethicone
Triglyceride

CHEMICAL AND PHYSICAL PROPERTIES (NOT PART OF SPECIFICATIONS)

Form	liquid
HLB value	approx. 5

APPLICATION

ABIL® EM 90 is a non-ionic W/O emulsifier which is based on silicone.

The high emulsion stabilizing potential of ABIL® EM 90 is caused by the polymeric and polyfunctional structure.

- ABIL® EM 90 is suitable for the formulation of W/O creams and lotions.

- Substances which can be processed include not only paraffin oils, which from the emulsion-technological point of view provide relatively few problems, but also fatty acid esters of short- and long-chain alcohols, silicone derivatives and vegetable triglycerides which are known to be difficult to emulsify.
- The amount used, referred to the emulsion, is only 1.5 – 2.5%. In formulations with a high water content the additional use of about 1% ISOLAN® GI 34 is recommended to improve the application properties.
- Consistency-providing or emulsion-stabilizing waxes are required only in amounts up to 2%. Amongst others, hydrogenated castor oil in combination with high-melting hydrocarbon waxes or beeswax are suitable.
- The optimum range for the content of oil phase is between 22 and 35%.
- Substances with specific properties, such as UV filters, plant extracts, moisturizers and antiperspirants, are well tolerated by the emulsion.
- The creams and lotions are distinguished by high stability towards heat and freezing stress; stability between -25 °C and +60 °C is attainable. ABIL® EM 90 can be used for the preparation of multiple emulsions of the type W/O/W and O/W/O. [P. Hameyer, K.R. Jenni, Emulsifiers for Multiple Emulsions – Optimization of stability by constitution and Molecular Weight, Parfümerie und Kosmetik 12 (1994), 842 – 850]
- Emulsions in which the oil phase contains predominantly substances with good spreading properties have good rub-out characteristics and are rapidly absorbed by the skin. On the other hand, creams which contain only highly viscous oils as emulsifiable substances, e. g. vegetable triglycerides, have a "heavy" action, whilst maintaining pleasant application properties.

INFLUENCE ON THE VISCOSITY OF THE EMULSION

The viscosity of the W/O emulsions based on ABIL® EM 90 can be regulated via three variables.

1. Viscosity of the oil phase

The viscosity of the external phase correlates directly with the viscosity of the emulsion. This means that it increases when low viscosity oils in a formulation are replaced by more viscous oils or when waxes are added; the latter show distinct effects even at 0.5 – 1.0%.

2. Phase ratio

In emulsions with predominant content of dispersed phase – as in the cosmetic W/O preparations – the viscosity increases significantly when the proportion of the dispersed phase is increased. The reason for this is the interaction between the dispersed water droplets which becomes stronger with increasing packing density. Therefore, ABIL® EM 90 creams on average have lower oil contents than lotions.


Depending on the emulsifiable substances, creams can be prepared about 20 – 33% of oil phase, lotions about 25 – 35%.

A given oil phase can be processed to form stable viscous liquid or cream-like emulsions when its proportion is in the optimum range for stability of the emulsion. E. g. decyl oleate forms creams with 19 – 21% of oil phase and lotions with 22 – 25% (see graph).

Outside these limits the stability of the emulsion is reduced because the dispersed water phase is too high or the viscosity of the emulsion too low. This working range is displaced to higher percentages when oils or oil/wax mixtures with higher viscosity are used.

3. Degree of dispersion

An additional parameter having an influence on viscosity is the degree of dispersion, which, however, should not be used for regulation of viscosity due to its effect on the stability of the emulsion. The viscosity increases when, due to mechanical processing, the diameter of the droplets is reduced and the specific boundary area between the phases is thereby increased. For this reason cream formulations are still liquid in the pre-emulsion state because of their coarse degree of dispersion.



PREPARATION

A pre-requisite is the careful adjustment of the formulation (phase ratio, viscosity of the oil phase) and optimum emulsification. The particle size for creams which are stable over a long period of time is below 1 µm, for lotions approx. 2 – 4 µm. More coarsely dispersed emulsions tend to separate.

Thorough, but not too intensive homogenization is required. Extreme energy input frequently causes the formation of highly viscous, metastable secondary structures which break down on storage. Under such conditions lotions can transiently assume cream-like consistency, e. g. by several passages through a colloid mill.

Optimum manufacturing conditions correspond to the principles of normal production processes for W/O emulsions. The water phase is incorporated slowly into the oil phase which contains the emulsifier while stirring intensively. The coarsely dispersed pre-emulsion is then homogenized. The final homogenization should be performed below 30 °C.

The temperature program is variable and can take the form of:

- hot/hot procedure (H/H)
- hot/cold procedure (H/C)

In addition to the traditional hot/hot procedure (both phases 80 – 90 °C) the hot/cold procedure can be used. It is characterized by incorporation of the cold water phase (15 – 30 °C) into the hot oil phase which significantly shortens the time of manufacture. Homogenization should be carried out below 30 °C in order to ensure that the waxes are largely recrystallized.

The decisive criterion for production is the viscosity. Mechanical processing is discontinued when the viscosity is equal to that of the standard emulsion developed and tested in the laboratory.

EMULSIFYING MACHINES

Stirring equipment or planetary mixers with high sheering force are very suitable for the manufacture of creams and lotions on the laboratory and production scale, provided that they guarantee uniform work-up of the emulsion. Machines predominately used in the cosmetic industry, which are equipped with stirrer, stripper and rotor-stator homogenizer, fulfill all requirements for optimum emulsification. However, utilization of their maximum capacity may result in over-emulsification. High-pressure emulsifiers may cause problems because of the danger of over-emulsification and liberation of water due to cavitation.

RECOMMENDED USAGE CONCENTRATION

1.5 – 2.5% of ABIL® EM 90

PACKAGING

180 kg drum

HAZARDOUS GOODS CLASSIFICATION

Information concerning

- classification and labeling according to regulations for transport and for dangerous substances
- protective measures for storage and handling
- measures in case of accidents and fires
- toxicity and ecological effects

is given in our material safety data sheets.

GUIDELINE FORMULATIONS

W/O Extra Moisture Cream BR 3/16-1

Phase A

ABIL® EM 90	2.00%
Mineral Oil (30 mPas)	17.00%
Hydrogenated Castor Oil	0.40%
Microcrystalline Wax (Paracera W 80, Paramelt B.V.)	0.60%

Phase B

Sodium Chloride	0.50%
Water	58.80%

Phase C

Urea	10.00%
Water	10.00%

Phase D

Phenoxyethanol, Ethylhexylglycerin (Euxyl PE 9010, Schülke&Mayr GmbH)	0.70%
--------------------------------------------------------------------------	-------

Preparation:

1. Heat phase A to approx. 80 °C.
2. Add phase B (80 °C or room temperature) slowly while stirring.
3. Homogenize for a short time.
Cool with gentle stirring. Add phase C and D at approx. 35 °C.
4. Homogenize again below 30 °C.

W/O Silk Body Milk SZ 10/14-1

Phase A

ABIL® EM 90	2.00%
Beeswax	0.50%
Hydrogenated Castor Oil	0.50%
Mineral Oil (30 mPas)	10.50%
TEGOSOFT® DC (Decyl Cocoate)	8.00%
Cyclomethicone	6.00%
Tocopheryl Acetate	0.50%

Phase B

Sodium Chloride	0.50%
Glycerin	3.00%
Water	62.80%

Phase C

Alcohol	5.00%
---------	-------

Phase D

Phenoxyethanol, Ethylhexylglycerin (Euxyl PE 9010, Schülke&Mayr GmbH)	0.70%
--------------------------------------------------------------------------	-------

Preparation:

1. Heat phase A to approx. 80 °C.
2. Add phase B (80 °C or room temperature) slowly while stirring.
3. Homogenize for a short time.
4. Cool with gentle stirring. Add phase C and D below 40 °C.
5. Homogenize again below 30 °C.

W/O Antiperspirant Roll-on H 13/09-27

Phase A

ABIL® EM 90	1.50%
ABIL® EM 97 S (Bis PEG/PPG-14/14 Dimethicone; Dimethicone)	1.50%
TEGOSOFT® DEC (Diethylhexyl Carbonate)	10.60%
Dimethicone (5 mPas)	10.60%
Silica (Aerosil R 812, Evonik Industries)	0.30%
Zinc Stearate	0.50%

Phase B

Water	51.30%
Glycerin	3.00%
Sodium Chloride	0.70%
Aluminum Chlorohydrate (Reach 501 L, Summit Reheis)	20.00%

Phase Z

Preservative, Perfume	q.s.
-----------------------	------

Preparation:

1. Disperse Zinc Stearate with ultra-turrax in emulsifiers and oils of phase A.
2. Disperse Aerosil R 812 in phase A.
3. Prepare phase B separately and add slowly while stirring to phase A.
4. Homogenize.

Disclaimer

This information and all further technical advice is based on our present knowledge and experience. However, it implies no liability or other legal responsibility on our part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments.

The customer is not released from the obligation to conduct careful inspection and testing of incoming goods. Performance of the product described herein should be verified by testing, which should be carried out only by qualified experts in the sole responsibility of a customer. Reference to trade names used by other companies is neither a recommendation, nor does it imply that similar products could not be used. (Status: April, 2008)

Evonik Nutrition & Care GmbH

Goldschmidtstraße 100

45127 Essen, Germany

P.O.BOX 45116 Essen

PHONE + 49 201 173-2546

FAX +49 201 173 712546

personal-care@evonik.com

www.evonik.com/personal-care

H 08/16